

# Arthritis of the Thumb

---

Arthritis is a condition that irritates or destroys a joint. Although there are several types of arthritis, the one that most often affects the joint at the [base](#) of the thumb (the basal joint) is osteoarthritis (degenerative or "wear-and-tear" arthritis).

## Description

---

Smooth cartilage covers the ends of the bones. It [enables](#) the bones to glide easily in the joint. Without it, bones rub against each other, causing friction and [damage](#) to the bones and the joint. Osteoarthritis occurs when the cartilage begins to wear away.

The joint at the base of the thumb, near the wrist and at the fleshy part of the thumb, enables the thumb to swivel, pivot, and pinch so that you can grip things in your hand.

Arthritis of the base of the thumb is more common in women than in men, and usually occurs after 40 years of age. Prior fractures or other injuries to the joint may increase the likelihood of developing this condition.

[Top of page](#)



## Symptoms

---

- Pain with activities that involve gripping or pinching, such as turning a key, opening a door, or snapping your fingers
- Swelling and tenderness at the base of the thumb
- An aching discomfort after prolonged use
- Loss of strength in gripping or pinching activities
- An [enlarged](#), "out-of-joint" appearance
- Development of a bony prominence or bump over the joint
- Limited motion

[Top of page](#)

## Doctor Examination

---

Your doctor will ask you about your symptoms, any prior [injury](#), pain patterns, or activities that aggravate the condition.

One of the tests used during the examination involves [holding](#) the joint firmly while moving the thumb. If pain or a gritty feeling results, or if a grinding sound (crepitus) can be heard, the bones are rubbing directly against each other.

An x-ray may show deterioration of the joint as well as any [bone spurs](#) or calcium deposits that have developed.

Many people with arthritis at the base of the thumb also have symptoms of [carpal tunnel syndrome](#), so your doctor may check for that, as well.

[Top of page](#)

## Treatment

---

## ***Nonsurgical Treatment***

In its early stages, arthritis at the base of the thumb will respond to [nonsurgical treatment](#).

- Ice the joint for 5 to 15 minutes several times a day.
- Take an anti-inflammatory [medication](#), such as aspirin or ibuprofen, to help reduce inflammation and swelling.
- Wear a supportive splint to [limit](#) the movement of your thumb, and allow the joint to rest and heal. The splint may protect both the wrist and the thumb. It may be worn [overnight](#) or intermittently during the day.

Because arthritis is a progressive, [degenerative disease](#), the condition may worsen over time. The next phase in treatment involves injecting a steroid solution injection directly into the joint. This will usually provide relief for several months. However, these injections cannot be repeated indefinitely.

## ***Surgical Treatment***

When nonsurgical treatment is no longer effective, [surgery](#) is an option. The operation can be performed on an outpatient basis, and several different procedures can be used.

One option involves fusing the bones of the joint together. This, however, will limit movement.

Another option is to remove part of the joint and reconstruct it using either a tendon graft or an artificial [substance](#).

You and your [physician](#) will discuss the options and select the one that is best for you.

[Top of page](#)

## **Rehabilitation**

---

After surgery, you will have to wear a cast for 4 to 8 weeks, depending on which procedure is used. A [rehabilitation program](#), often involving a physical therapist, helps you regain movement and strength in the hand. You may feel some discomfort during the initial stages of the rehabilitation program, but this will diminish over time. Full recovery from surgery takes several months. Most patients are able to resume normal activities.

[Top of page](#)

Last reviewed: December 2013

*AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS ["Find an Orthopaedist"](#) program on this web site.*